VOL. 517 JOURNAL OF CHROMATOGRAPHY SEPTEMBER 26, 1990

CONTENTS

HONOUR VOLUME ON THE OCCASION OF THE 60TH BIRTHDAY OF J. CALVIN GIDDINGS

Preface
by K. Caldwell 1

Plate height theory for compressible mobile phase fluids and its application to gas, liquid and supercritical fluid chromatography
by D. P. Poe (Duluth, MN, U.S.A.) and D. E. Martire (Washington, DC, U.S.A.) . . . 3

Dispersion in the interstitial space of packed columns
by P. Magnico and M. Martin (Paris, France) 31

Comparison of methods for the determination of the polarity and selectivity of stationary phases in gas chromatography from a thermodynamic point of view
by R. V. Golovnya and B. M. Polanuer (Moscow, U.S.S.R.) 51

Cryogenic-focusing, ohmically heated on-column trap for capillary gas chromatography
by S. R. Springston (Upton, NY, U.S.A.) 67

Enhanced possibilities for identification using series-coupled capillary gas chromatographic columns. II. Retention indices as an identification tool in selectivity tuning
by T. Maurer, W. Engewald and A. Steinborn (Leipzig, G.D.R.) 77

Oxygen enhancement of thermal electron capture in a non-radioactive discharge source for a quadrupole mass spectrometer
by W. E. Wentworth, E. D. D'sa, C. F. Batten and E. C. M. Chen (Houston, TX, U.S.A.) 87

Comparison of cuticular hydrocarbon profiles of fire ants *Solenopsis richteri* from the same colony, using capillary column gas chromatography with pattern recognition
by J. H. Brill and W. Bertsch (Tuscaloosa, AL, U.S.A.) 95

Empirical scheme for the classification of gas chromatographic stationary phases based on solvatochromic linear solvation energy relationships
by J. Li, A. J. Dallas and P. W. Carr (Minneapolis, MN, U.S.A.) 103

Capillary gas chromatography with graphitized carbon black
by F. Bruner, G. Crescentini, F. Mangani and L. Lattanzi (Urbino, Italy) 123

Pesticide analysis by gas chromatography with a novel atomic emission detector
by P. L. Wylie (Avondale, PA, U.S.A.) and R. Oguchi (Tokyo, Japan) 131

Turbulent flow in cappillary gas chromatography—evaluation of a theoretical concept by Golay
by A. J. van Es, J. A. Rijks and C. A. Cramers (Eindhoven, The Netherlands) and M. J. E. Golay (Norwalk, CT, U.S.A.) 143

Use of chromatographic models for computerized optimization of coupling-point pressure in dual-column gas chromatography
by T. Hevesi and J. Krupčík (Bratislava, Czechoslovakia) and P. Sandra (Ghent, Belgium) 161

Study of rapid reaction kinetics by computerized gas chromatography with stroboscopic sampling
by E, Küllik and M. Kaljurand (Estonia, U.S.S.R.) 175

High-performance liquid chromatographic study of ligand-exchange reactions of fluorinated metal β-diketone chelates
by T. Wang and P. C. Uden (Amherst, MA, U.S.A.) 185

Silicas chemically bonded with multidentate phenyl groups as stationary phases in reversed-phase liquid chromatography used for non-planarity recognition of polycyclic aromatic hydrocarbons
by K. Jinno, K. Yamamoto, H. Nagashima, T. Ueda and K. Itoh (Toyohashi, Japan) . . 193

Gel permeation chromatography–Fourier transform infrared study of some synthetic polymers. II. Instrumentation for the characterization of polyethylene
by K. Nishikida (Yokohama, Japan), T. Housaki (Chiba, Japan) and M. Morimoto and T. Kinoshita (Yokohama, Japan) 209

Isolation and purification of lecithin by preparative high-performance liquid chromatography
by J. V. Amari and P. R. Brown (Kingston, RI, U.S.A.), C. M. Grill (Wakefield, RI, U.S.A.) and J. G. Turcotte (Kingston, RI, U.S.A.) 219

Optimization of the experimental conditions in preparative liquid chromatography with touching bands
by S. Golshan-Shirazi and G. Guiochon (Oak Ridge, TN, U.S.A.) 229

Evaluation of the stability of polymer-coated silica-based packing materials for high-performance liquid chromatography
by T. Takeuchi, W. Hu, H. Haraguchi and D. Ishii (Nagoya, Japan) 257

Magnetic resonance imaging in reversed-phase liquid chromatography
by M. Ilg, J. Maier-Rosenkranz, W. Müller and E. Bayer 263

Non-porous polybutadiene-coated silicas as stationary phases in reversed-phase chromatography
by M. Hanson and K. K. Unger (Mainz, F.R.G.) and G. Schomburg (Mülheim, F.R.G.) . 269

Competition between phenylalanine and acetic acid in a chromatographic column as indicated by their adsorption isotherms
by S. Levin and S. Abu-Lafi (Jerusalem, Israel) 285

Design of liquid chromatography capillary columns
by R. P. W. Scott (Washington, DC, U.S.A.) 297

Reversed-phase ion-pair chromatography with indirect photometric detection of inorganic anions from residues of low explosives
by D. Woolfson-Bartfeld, E. Grushka, S. Abramovich-Bar, S. Levy and Y. Bamberg (Jerusalem, Israel) 305

High-performance liquid chromatography of amino acids, peptides and proteins. C. Characterisation of coulombic interactive regions on hen lysozyme by high-performance liquid anion-exchange chromatography and computer graphic analysis
by A. N. Hodder (Clayton, Australia), K. J. Machin (Fitzroy, Australia) and M. I. Aguilar and M. T. W. Hearn (Clayton, Australia) 317

Optimization study of octane-in-water emulsions by sedimentation field-flow fractionation
by M. E. Hansen and D. C. Short (Cincinnati, OH, U.S.A.) 333

Potential-barrier field-flow fractionation, a versatile new separation method
by A. Koliadima and G. Karaiskakis (Patras, Greece) 345

Particle characterization in centrifugal fields. Comparison between ultracentrifugation and sedimentation field-flow fractionation
by J. Li and K. D. Caldwell (Salt Lake City, UT, U.S.A.) and W. Mächtle (Ludwigshafen, F.R.G.) 361

Retention effects in thermal field-flow fractionation
by J. J. Kirkland, L. S. Boone and W. W. Yau (Wilmington, DE, U.S.A.) 377

Sedimentation field-flow fractionation for pigment quality assessment
by L. Koch, T. Koch and H. M. Widmer (Basle, Switzerland) 395

Characterization of polymers by thermal field-flow fractionation
by M. E. Schimpf (Boise, ID, U.S.A.) 405

Evaluation of pinched inlet channel for stopless flow injection in steric field-flow fractionation
by M. H. Moon, M. N. Myers and J. C. Giddings (Salt Lake City, UT, U.S.A.) 423

Use of field-flow fractionation to study pollutant–colloid interactions
by R. Beckett, D. M. Hotchin and B. T. Hart (Caulfield East, Australia) 435

Quantitative correlations between solute molecular structure and solubility in supercritical fluids
by J. W. King and J. P. Friedrich (Peoria, IL, U.S.A.) 449

Use of chromatographic retention measurements to obtain solubilities in a liquid or supercritical fluid mobile phase
by K. D. Bartle, A. A. Clifford, S. A. Jafar, J. P. Kithinji and G. F. Shilstone (Leeds, U.K.) 459

Element-selective detection after supercritical fluid chromatography by means of a Surfatron plasma in the near-infrared spectral region
by D. R. Luffer and M. Novotny (Bloomington, IN, U.S.A.) 477

Use of sulfur chemiluminescence detection after supercritical fluid chromatography
by H.-C. K. Chang and L. T. Taylor (Blacksburg, VA, U.S.A.) 491

Static coating of 5 to 50 μm I.D. capillary columns for open tubular column chromatography
by S. R. Sumpter, C. L. Woolley, E. C. Huang, K. E. Markides and M. L. Lee (Provo, UT, U.S.A.) 503

Influence of thermal variation of diffusion coefficient on non-equilibrium plate height in capillary zone electrophoresis
by J. M. Davis (Carbondale, IL, U.S.A.) 521

Micellar electrokinetic chromatography employing sodium alkyl sulfates and Brij 35®
by H. T. Rasmussen, L. K. Goebel and H. M. McNair (Blacksburg, VA, U.S.A.) 549

Electrically driven open-tubular liquid chromatography
by G. J. M. Bruin, P. P. H. Tock, J. C. Kraak and H. Poppe (Amsterdam, The Netherlands) 557

High-pressure and supercritical capillary electrophoresis
by C. R. Yonker and R. D. Smith (Richland, WA, U.S.A.) 573

Author Index 583

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)